

Manage postpartum hemorrhage with the Bakri® Postpartum Balloon with Rapid Instillation Components

Postpartum hemorrhage guidelines

- California Maternal Quality Care Collaborative. Obstetric hemorrhage emergency management plan: table chart. CMQCC Web site. <https://www.cmqcc.org/resource/3304/download>. Published March 24, 2015. Accessed October 5, 2020.
- Committee on Practice Bulletins–Obstetrics. Practice bulletin no. 183: postpartum hemorrhage. *Obstet Gynecol*. 2017;130(4):e168-e186.
- Prevention and management of postpartum haemorrhage: Green-top guideline no. 52. *BJOG*. 2017;124(5):e106–e149.
- Sentilhes L, Vayssière C, Deneux-Tharoux C, et al. Postpartum hemorrhage: guidelines for clinical practice from the French College of Gynaecologists and Obstetricians (CNGOF): in collaboration with the French Society of Anesthesiology and Intensive Care (SFAR). *Eur J Obstet Gynecol Reprod Biol*. 2016;198:12-21.
- Takeda S, Makino S, Takeda J, et al. Japanese clinical practice guide for critical obstetrical hemorrhage (2017 revision). *J Obstet Gynaecol Res*. 2017;43(10):1517-1521.
- World Health Organization. WHO recommendations: uterotonics for the prevention of postpartum haemorrhage. WHO Web site. <https://apps.who.int/iris/bitstream/handle/10665/277276/9789241550420-eng.pdf>. WHO Published 2018. Accessed November 17, 2020.

Systematic review and meta-analysis

- Suarez S, Conde-Agudelo A, Borovac-Pinheiro A, et al. Uterine balloon tamponade for the treatment of postpartum hemorrhage: a systematic review and meta-analysis. *Am J Obstet Gynecol*. 2020;222(4):293.e1-293.e52.

Multicenter study

- Vintejeux E, Ulrich D, Mousty E, et al. Success factors for Bakri™ balloon usage secondary to uterine atony: a retrospective, multicentre study. *Aust N Z J Obstet Gynaecol*. 2015;55(6):572-577.
- Wang D, Xu S, Qiu X, et al. Early usage of Bakri postpartum balloon in the management of postpartum hemorrhage: a large prospective, observational multicenter clinical study in South China. *J Perinat Med*. 2018;46(6):649-656.

Bakri vs. invasive procedure

- Gauchotte E, De La Torre M, Perdriolle-Galet E, et al. Impact of uterine balloon tamponade on the use of invasive procedures in severe postpartum hemorrhage. *Acta Obstet Gynecol Scand*. 2017;96(7):877-882.

- Laas E, Bui C, Popowski T, et al. Trends in the rate of invasive procedures after the addition of the intrauterine tamponade test to a protocol for management of severe postpartum hemorrhage. *Am J Obstet Gynecol*. 2012;207(4):281.e1-281.e7. doi:10.1016/j.ajog.2012.08.028.
- Lo A, St Marie P, Yadav P, et al. The impact of Bakri balloon tamponade on the rate of postpartum hysterectomy for uterine atony. *J Matern Fetal Neonatal Med*. 2017;30(10):1163-1166.
- Maher MA, Abdelaziz A. Comparison between two management protocols for postpartum hemorrhage during cesarean section in placenta previa: balloon protocol versus non-balloon protocol. *J Obstet Gynaecol Res*. 2017;43(3):447-455.
- Revert M, Rozenberg P, Cottenet J, et al. Intrauterine balloon tamponade for severe postpartum hemorrhage. *Obstet Gynecol*. 2018;131(1):143-149.

Placenta previa, accrete, and increta

- Lin B, Zhou B, Chen J, et al. Prophylactic application of Bakri balloon tamponade versus uterine gauze packing during cesarean section in patients with placenta previa. *J Int Med Res*. 2020;48(3):300060520910049.
- Pala Ş, Atilgan R, Başpınar M, et al. Comparison of results of Bakri balloon tamponade and caesarean hysterectomy in management of placenta accreta and increta: a retrospective study. *J Obstet Gynaecol*. 2018;38(2):194-199.
- Soyama H, Miyamoto M, Ishibashi H, et al. Analysis of prophylactic Bakri balloon tamponade failure in patients with placenta previa. *Taiwan J Obstet Gynecol*. 2019;58(1):159-163.
- Soyama H, Miyamoto M, Sasa H, et al. Effect of routine rapid insertion of Bakri balloon tamponade on reducing hemorrhage from placenta previa during and after cesarean section. *Arch Gynecol Obstet*. 2017;296(3):469-474.
- Tahaoglu AE, Balsak D, Erdogdu E, et al. Bakri balloon placement effectively treats uterine atony and placenta previa. *Biotechnol Biotechnol Equip*. 2017;31(4):795-799.

Tamponade test

- Georgiou C. Intraluminal pressure readings during the establishment of a positive 'tamponade test' in the management of postpartum haemorrhage. *BJOG*. 2010;117(3):295-303.
- Kong CW, To WWK. Intraluminal pressure of uterine balloon tamponade in the management of severe postpartum hemorrhage. *J Obstet Gynaecol Res*. 2018;44(5):914-921.
- Li H, Bai TY, Bai L, et al. Intraluminal pressure difference of Bakri balloon in the positive tamponade test during the management of severe post-partum hemorrhage. *Biomed J Sci Tech Res*. 2019;15(4):11524-11527.

General

- Brown H, Okeyo S, Mabeya H, et al. The Bakri tamponade balloon as an adjunct treatment for refractory postpartum hemorrhage. *Int J Gynaecol Obstet*. 2016;135(3):276-280.
- Kong CW, To WW. Prognostic factors for the use of intrauterine balloon tamponade in the management of severe postpartum hemorrhage. *Int J Gynaecol Obstet*. 2018;142(1):48-53.
- Mathur M, Ng QJ, Tagore S. Use of Bakri balloon tamponade (BBT) for conservative management of postpartum haemorrhage: a tertiary referral centre case series. *J Obstet Gynaecol*. 2018;38(1):66-70.